

## **Characterization of Plant Species Diversity and Abundance to Inform Effective Rehabilitation of Cherangany Hills Forest**

Nereoh Leley<sup>1</sup>, \* Jared Amwatta Mullah<sup>2</sup>, Michael Macharia<sup>3</sup>, Boaz Ngonga<sup>1</sup> and Jackline Atieno<sup>4</sup>

<sup>1</sup> Kenya Forestry Research Institute, Rift Valley Eco-Region Research Program-Londiani

<sup>2</sup> Kenya Forestry Research Institute, Lake Victoria Basin Eco-Region Research Program-Maseno

<sup>3</sup> Kenya Forestry Research Institute, Central Highlands Eco-Region Research Program-Muguga

<sup>4</sup> Egerton University, Department of Natural Resource Management-Njoro

\*Email: [nereoh24@gmail.com](mailto:nereoh24@gmail.com):

### **Abstract**

Cherangany Hills Forest contains high levels of tree species diversity and endemism and is in critical need of conservation and restoration. Little is known about natural regeneration dynamics of isolated forest fragments forming the Cherangany, or the potential for human action to aid recovery of lost structures and functions following deforestation/degradation. Here, data and analyses are presented from a series of rapid ecological surveys conducted in the main forest fragments of highly impacted, low impacted and no impact forest mosaics. A total of 337 plant species were recorded which is an improvement from a total of 302 plant species recorded in the last survey done in early 1990s. Embobut forest block which in the highly impacted category had the lowest regeneration potential in contrast to Kipkunur block, low impacted fragment which had the highest. Study results indicated that continued human disturbances has led to diminishing density of key commercial hard timber tree species like *Hagenia abyssinica*, *Olea welwitschii*, *Warbugia ugandensis*, and *Juniperus procera*. The low density of both mature trees and seedlings in degraded sites may undermine the long-term viability of these key species. Selection of potential restoration tree species for each fragment should consider the density of individual tree species with priority given to species with low density for rehabilitation. Furthermore, analysis of illegally harvested trees highlighted the important role of indigenous tree species as a source of ecosystem services to local people; an important consideration for forest conservation and rehabilitation planning for the ecosystem.

**Keywords:** Water Towers, Degradation, Regeneration, Management response